

Critical Casimir Effect in ^3He - ^4He films

A. Maciołek¹ and S. Dietrich^{2,3}

¹*Institute of Physical Chemistry, Polish Academy of Sciences,*

Kasprzaka 44/52, PL-01-224 Warsaw, Poland

²*Max-Planck-Institut für Metallforschung,*

Heisenbergstr. 3, D-70569 Stuttgart, Germany

³*Institut für Theoretische und Angewandte Physik,*

Universität Stuttgart, Pfaffenwaldring 57, D-70569 Stuttgart, Germany

(Dated: February 2, 2008)

Abstract

Universal aspects of the thermodynamic Casimir effect in wetting films of ^3He - ^4He mixtures near their bulk tricritical point are studied within suitable models serving as representatives of the corresponding universality class. The effective forces between the boundaries of such films arising from the confinement are calculated along isotherms at several fixed concentrations of ^3He . Nonsymmetric boundary conditions impose nontrivial concentration profiles leading to repulsive Casimir forces which exhibit a rich behavior of the crossover between the tricritical point and the line of critical points. The theoretical results agree with published experimental data and emphasize the importance of logarithmic corrections.

PACS numbers: 05.70.Jk, 64.60.Fr, 64.60.Kw, 67.40.Kh, 68.15.+e

Finite-size contributions to the free energy of a fluid confined between two surfaces at a distance L give rise to an effective force between them. Theory predicts that at the bulk critical point T_c of such a system this force becomes long-ranged as a result of critical fluctuations of the corresponding ordering degrees of freedom. This is analogous to the well-known Casimir effect in electromagnetism. This so-called critical Casimir force f_C per unit area and in units of $k_B T_c$ can be expressed in terms of universal scaling functions [1].

Only recently, sophisticated wetting experiments have provided detailed quantitative data for critical Casimir forces in various systems [2, 3, 4, 5, 6]. In the case of ^4He wetting films near the superfluid transition, these experimental studies support quantitatively theoretical predictions for f_C ($T \geq T_c$) corresponding to the universality class of the XY model [7]. For the case of ^3He - ^4He films near the bulk tricritical point some theoretical predictions are available [7], but those do not apply for the boundary conditions relevant for recent wetting experiments performed in these systems [4, 5]. However, the shape of the scaling function of the Casimir force depends *sensitively* on the type of boundary conditions (BC) and thus on the surface universality classes to which the confining surfaces belong [8]. The experiments of Ref. [4] report a *repulsive* f_C around the tricritical point which suggests *nonsymmetric* BC for the superfluid order parameter (SOP). This is opposite to the case of pure ^4He wetting films near the λ -point where f_C was found to be *attractive* [2, 7]. For the latter system the BC seem to be very well approximated by *symmetric* Dirichlet-Dirichlet BC (O, O) forming the so-called ordinary (O) surface universality class because the quantum-mechanical wave function describing the superfluid state vanishes at both interfaces [1, 2]. The type of BC for ^3He - ^4He wetting films is not clear from the outset because a ^4He -rich layer forms near the substrate-fluid interface, which may become superfluid already above the bulk λ -line [9] whereas ^3He has a preference for the fluid-vapor interface. Thus the two interfaces impose a nontrivial concentration profile which in turn couples to the SOP. This leads to the hypothesis that the concentration profile induces effectively nonsymmetric ($O, +$) BC for the SOP, i.e., Dirichlet boundary conditions at the fluid-vapor interface and symmetry-breaking ($+$) BC at the substrate-fluid interface (also known as the so-called extraordinary or normal universality class [8]). For the present tricritical behavior the upper critical dimension d^* equals 3. In this case theory predicts that for three-dimensional systems the asymptotic tricritical thermodynamic functions exhibit power laws with critical exponents taking their classical values. However, logarithmic corrections to the mean-field (MF) behavior are expected

under experimental conditions [10].

Here we consider two complementary approaches. Field-theoretical methods and renormalization-group (RG) analyses are used to derive universal properties of the Casimir force at the tricritical point and the form of logarithmic corrections. However, these methods do not lend themselves for systematic studies of f_C along all thermodynamic paths followed in the aforementioned experiments. In order to be able to interpret the rich variation of f_C extracted from the capacity measurements in Ref. [4], to understand the emergence of the actual BC and, moreover, to predict the behavior of f_C in the crossover region between the tricritical and the critical points, we employ the vectorized Blume-Emery-Griffiths model (VBEG) [12] as a representative of the same universality class as the actual physical system. This lattice model is extended to the film geometry and treated within mean field theory (MFT).

First we derive the leading asymptotic behavior of f_C at tricriticality for $(O, +)$ BC. To this end we consider the standard Ginzburg-Landau (GL) Hamiltonian for an $O(n)$ -symmetric tricritical system ($T = T_t$) in a film geometry:

$$\mathcal{H}[\Phi] = \int d^{d-1}x \int_0^L dz \left\{ \frac{1}{2}(\nabla\Phi)^2 + \frac{u}{6!}(\Phi^2)^3 \right\} \quad (1)$$

where L is the film thickness, Φ is the n -component OP and z is the distance between the confining surfaces; u is a bare coupling constant. In a film geometry $f_C \equiv -(\partial f^{ex}/\partial L) = \langle \mathcal{T}_{zz} \rangle$ is given by the stress tensor component $\mathcal{T}_{zz}$ [1], where $f^{ex}(L) \equiv (f - f_b)L$, f is the total free energy per unit area and per $k_B T_t$ and f_b is the bulk contribution. The stress tensor is given by [1] $\mathcal{T}_{ij} = \partial_i \Phi \cdot \partial_j \Phi - \delta_{ij} \mathcal{L} - (d-2)/(4(d-1))(\partial_i \partial_j - \delta_{ij} \nabla^2) \Phi^2$, where $\mathcal{L}$ is the integrand of (1). We take $\Phi = (m(z), 0, \dots, 0)$. Determination of the tricritical Casimir force starts from the Euler-Lagrange equation for the OP profile: $m''(z) = (u/120)m^5(z)$ with $(O, +)$ BC, i.e., $m(0) = 0$ and $m(L) = +\infty$. In this case the spatially constant $\langle \mathcal{T}_{zz} \rangle$ can be expressed as $(1/2)(m'(0))^2$. With the scaling ansatz $m(z) = (u/360)^{-1/4} L^{-1/2} \varphi(z/L)$ and $\langle \mathcal{T}_{zz} \rangle = (90/u)^{1/2} L^{-3} \Theta$, and after integrating directly the first integral of the Euler-Lagrange equation one obtains $\Theta^{1/3} = \int_0^\infty dp / \sqrt{1+p^6} \simeq 1.40218$ by implementing the above BC. Eventually, in units of $k_B T_t$ the MFT result for the tricritical Casimir force f_C^t in the case of $(O, +)$ BC is $f_C^t = 2.7568(4) (90/u)^{1/2} L^{-3}$. Note that within MFT the parameter $u > 0$ remains undetermined. Its value follows from using standard RG arguments. In $d = 3 - \epsilon$ the above MFT result yields the leading contribution in an ϵ -expansion. After

removing the uv singularity via renormalization the asymptotic scaling behaviour of f_C^t follows from substituting u by the appropriate fixed-point value $u^* \propto \epsilon$. At $d = d^*$, and under spatial rescaling by a factor ℓ , u flows to its RG fixed point value $u^* = 0$ according to $\bar{u}(\ell) = (240\pi^2)/((3n+22)|\ln \ell|)$ [11]. With the rescaling factor $\ell = l_0/L$, where l_0 is a microscopic length scale of the order of a few Å, this yields a logarithmic correction to the power law L -dependence of the tricritical Casimir force:

$$f_C^t \simeq 0.54(3n+22)^{1/2}(\ln L/l_0)^{1/2}L^{-3}. \quad (2)$$

Gaussian fluctuations give contributions of at least $O(u^0)$ which are therefore subdominant. We compare Eq. (2) for $n = 2$ with the data obtained by Garcia and Chan [4] for their experimental value $L/l_0 \approx 520 \text{ Å}/1.3 \text{ Å}$. This gives $\vartheta_t \equiv f_C^t L^3 \approx 6.96$ in a good agreement with $\vartheta_t^{exp} = 8.4 \pm 1.7$, which suggests that this experiment maybe the first to have verified implicitly the existence of logarithmic corrections near the tricritical point. However, in order to extract the actual value of the *universal* Casimir amplitude (i.e., the numerical prefactor in Eq. (2)) the experimental data require a reanalysis based on a functional form given by Eq. (2).

Now we turn to the VBEG model and consider a $d = 3$ simple cubic lattice consisting of L parallel lattice layers at spacing a . Each layer has A sites, labeled $i, j, \dots$ and associated with an occupation variable $t_i = 0, 1$ and a phase θ_i ($0 \leq \theta_i < 2\pi$) which mimics the phase of the ^4He wave function. A ^3He (^4He) atom at site i corresponds to $t_i = 0(1)$. The Hamiltonian is given by

$$\mathcal{H} = -J \sum_{\langle ij \rangle} t_i t_j \cos(\theta_i - \theta_j) - K \sum_{\langle ij \rangle} t_i t_j + \sum_i \Delta_i t_i \quad (3)$$

where the first two sums run over nearest-neighbor pairs and the last one is over all lattice sites. The field Δ_i is related to the chemical potentials of the two components of the mixture. $\Delta_i = \Delta_1$ and $\Delta_i = \Delta_2$ on the left and right surface layer, respectively, and $\Delta_i = \Delta$ otherwise. The differences $\Delta_i - \Delta$, $i = 1, 2$, are a measure of the relative preferences of ^4He atoms for the two surfaces such that $\Delta_1 < \Delta$ corresponds to the preference of ^4He atoms for the solid substrate. At the opposite surface with the vapor we choose $\Delta_2 = \Delta_t$, the tricritical bulk value. Near the tricritical point this choice is consistent with the assumption made in Ref. [4] for the concentration profile across the wetting film, whereby at the interface with the vapor the ^3He mole fraction takes the bulk value. Bulk properties of this model were studied within MFT and by Monte Carlo simulations in $d = 3$ in [12]. The resulting bulk phase

diagram resembles that observed experimentally for ^3He - ^4He mixtures (see Fig. 1). For the film geometry we have solved this model within MFT. This yields a set of self-consistent equations for the OP in the l th layer, i.e., the concentration $Q_l = \langle t_l \rangle = 1 - X(l)$ of ^4He , and the SOP $\mathbf{M}_l = (m_l^{(1)}, m_l^{(2)})$ with $m_l^{(1)} = \langle t_l \cos \theta_l \rangle$ and $m_l^{(2)} = \langle t_l \sin \theta_l \rangle$ where t_l and θ_l denote the occupation number and the phase in the l th layer, respectively. In the absence of helicity one has $\mathbf{M}_l = (m_l^{(1)}, 0) \equiv (m_l, 0)$, $Q_l = I_0(\tilde{J}b_l) / (e^{-\tilde{K}a_l + \Delta_l/T} + I_0(\tilde{J}b_l))$, and $m_l = I_1(\tilde{J}b_l) / (e^{-\tilde{K}a_l + \Delta_l/T} + I_0(\tilde{J}b_l))$. $I_0(z)$ and $I_1(z)$ are modified Bessel functions and $\Delta_l = \Delta$ for $l \neq 1, L$; $\tilde{K} = qK/T$ and $\tilde{J} = qJ/T$, where $q = q_\perp + 2q'$ is the coordination number of the lattice. $q_\perp$ is the in-layer coordination number while each site (but not in the first and last layers) is connected to q' atoms in each adjacent layer. We have introduced $b_l \equiv m_{l-1} + q_\perp m_l + m_{l+1}$ for $l \neq 1, L$, $b_1 \equiv q_\perp m_1 + m_2$, and $b_L \equiv m_{L-1} + q_\perp m_L$ and analogously $a_l \equiv Q_{l-1} + q_\perp Q_l + Q_{l+1}$ for $l \neq 1, L$, $a_1 = q_\perp Q_1 + Q_2$, and $a_L = Q_{L-1} + q_\perp Q_L$. The coupled equations for Q_l and m_l are solved numerically; the acceptable solution minimizes the free energy. First, we have analyzed the semi-infinite system. Close to the λ -line we observe a higher ^4He concentration near the left surface, which induces a local superfluid ordering (see Fig. 2). By varying T and Δ one obtains a whole line of continuous surface transitions corresponding to the onset of the formation of a superfluid film near the wall; it meets the λ -line at the special transition point whose position depends on the value of Δ_1 (see Fig. 1). These findings are in agreement with the results of a Migdal-Kadanoff analysis [13]. In the film geometry the Casimir force is obtained by taking a finite difference after calculating f^{ex} for L_0 and $L_0 + 1$. Figure 3 summarizes our result for a film of width $L = 20$, $K/J = 0.5$, $\Delta_1/J = -3$, and $\Delta_2 = \Delta_t/J \simeq 0.61$. f_C is calculated along the thermodynamic paths indicated in Fig. 1. Below T_t , f_C is calculated along the coexistence line, infinitesimally on the superfluid branch of bulk coexistence. Our results are presented in terms of the scaling function $\vartheta \equiv L^d f_C$ as a function of the scaling variable $y \equiv tL^{1/\nu} = (L\xi_0/\xi)^{1/\nu}$, where $t = (T - T_t)/T_t$. ξ_0 is the amplitude of the correlation length ξ and $\nu = 1$. The surface transition does not leave a visible trace in the behavior of f_C . For ^3He concentration $X < X_t$, upon crossing the λ -line there is a steep variation associated with a break in slope, giving rise to the formation of shoulders which are similar to those observed experimentally [4]. For $X > X_t$, when T reaches the phase separation temperature f_C coincides with the curve common to all values of X . This occurs with a discontinuous first derivative. The aforementioned common curve exhibits a pronounced maximum below T_t at $y \simeq -0.74$ and

gradually decreases to zero for $y \rightarrow -\infty$. Below T_t , both the concentration and the SOP profiles corresponding to this common curve display an interface-like structure separating two domains of the coexisting bulk phases (see $t = -0.0625$ in Fig. 2). Features of f_C in this 'soft mode' phase can be attributed to purely interfacial effects, similarly to Ising-like films with asymmetric BC [14]. Beyond MFT a positive sign of the force can be regarded as a consequence of entropic repulsion [15]. The maximum of f_C is expected to occur at the temperature T for which the interfacial width $\sim \xi_b \sim L$, i.e., $y \simeq -1$ [14] which checks with Fig. 3 [16]. For $X \leq X_t - 0.05$ we observe a crossover to the critical superfluid behavior of pure ^4He and a gradual formation of a second, less pronounced local maximum located slightly below the λ -line. This local maximum decreases upon departure from X_t and finally disappears above the special transition S . This is expected, because above S the BC turn into the type (O, O) for which f_C vanishes within MFT. For lower T , f_C increases steeply upon approaching bulk coexistence revealing that interfacial effects associated with the 'soft mode' lead to a much stronger Casimir effect than near the critical λ -line. The qualitative features of ϑ extracted from the experimental data for $X \simeq X_t$ (see Fig. 5 in Ref. [4]) are very well captured by the present lattice model. Discrepancies can be attributed to fluctuation effects neglected in the present MFT VBEG approach: (i) The discontinuities of slopes as obtained within MFT upon crossing the λ -line are expected to be smeared out by fluctuations. (ii) The experimental scaling function ϑ does not vanish at low temperatures, which may be due to Goldstone modes in the superfluid phase. However, the possibility that this behavior is an artifact of an extreme change in the dielectric constant of the film cannot be excluded [17]. In the crossover regime to the critical behavior only few experimental data for the thickness of the wetting film are published. However, again the variation of film thicknesses agrees with our findings. In particular, one observes a rapid thickening of the films upon approaching the coexistence line; for some values of X a small maximum located near the λ -line is also visible (compare Fig. 3). A quantitative comparison is not possible because, for our choice of surface terms in the Hamiltonian, the fixed-point BC $(O, +)$ cannot be reached within the VBEG model. In order to be able to extract universal properties - which requires to reach fixed-point BC - it would be necessary to introduce a surface field which couples to the SOP so that the BC $(+)$ can be realized. Also MFT is not sufficient, even in $d = d^*$. A naive correction of ϑ by the logarithmic factor derived within the GL model will not give the proper universal behavior. Instead, renormalization group

schemes or Monte Carlo simulations have to be employed. Nonetheless, our MFT results for $X = X_t$, if matched at the tricritical point $y = 0$ and after adjusting the amplitude ξ_0 so that the positions of the maximum of the scaling function are the same (i.e., $\xi_0^{th}/a \approx 0.065$) reproduce very well the experimental curve (see Fig. 4), especially near the maximum where we expect interfacial effects to be dominant. This is consistent because the 'soft mode' phase does not depend on the details of the surface fields. Notice, that the experimental data nominally for $X = X_t$ more closely match the theory for $X = X_t - 0.01$. This raises the question as to whether the ^3He concentration in the film is shifted relative to the bulk one.

A.M. benefited from discussions with R. Garcia, M. Krech and S. Kondrat. This work was partially funded by KBN grant No.4 T09A 066 22.

-
- [1] M. Krech, *The Casimir Effect in Critical System* (World Scientific, Singapore, 1994); J. Phys. Condens. Matter **11**, R391 (1999).
 - [2] R. Garcia and M. H. W. Chan, Phys. Rev. Lett. **83**, 1187 (1999).
 - [3] A. Mukhopadhyay and B. M. Law, Phys. Rev. Lett. **83**, 772 (1999).
 - [4] R. Garcia and M. H. W. Chan, Phys. Rev. Lett. **88**, 086101 (2002).
 - [5] T. Ueno, S. Balibar, T. Mizusaki, F. Caupin, and E. Rolley, Phys. Rev. Lett. **90**, 116102 (2003).
 - [6] M. Fukuto, Y. F. Yano, and P. S. Pershan, Phys. Rev. Lett. **94**, 135702 (2005).
 - [7] M. Krech and S. Dietrich, Phys. Rev. Lett. **66**, 345 (1991); **67**, 1055 (1991); Phys. Rev. A **46**, 1886 (1992); **46** 1922 (1992).
 - [8] H. W. Diehl, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, London, 1986), Vol. 10, p.76.
 - [9] J.-P. Romagnan, J.-P. Laheurte, J. -C. Noiray, and W. F. Saam, J. Low Temp. Phys. **30**, 425 (1978).
 - [10] D. Lawrie and S. Sarbach, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, London, 1984), Vol. 9, p.2.
 - [11] E. Eisenriegler and H. W. Diehl, Phys. Rev. B, **37**, 5257 (1988).
 - [12] A. Maciołek, M. Krech, and S. Dietrich, Phys. Rev. E **69**, 036117 (2004); and references

therein.

- [13] A. Crisanti and L. Peliti, J. Phys. A: Math. Gen. **18** L543 (1985).
- [14] A. O. Parry and R. Evans, Phys. Rev. Lett. **64**, 439 (1990).
- [15] M. E. Fisher, J. Chem. Soc. Faraday Trans. II **82**, 1569 (1986).
- [16] The experimental data for f_C exhibit a maximum at $y \simeq -18$ does not relate to the condition $\xi_b \sim L$.
- [17] R. Garcia, private communication.




FIG. 1: Phase diagram for the VBEG model obtained within MFT for $K/J = 0.5$ and $\Delta_1/J = -3$ exhibiting the bulk λ -line $T_s(X)$ of continuous superfluid transitions (long-dashed line), the phase separation curves (solid lines), the tricritical point $A = (T/T_s(0) = 2/3, X = 1/3)$, and the surface transition line (short-dashed line) which merges with the bulk λ -line at the special transition point $S = (T/T_s(0) \simeq 0.759, X \simeq 0.241)$. Vertical lines represent thermodynamic paths along which the Casimir force has been calculated (see Fig. 3). $\blacklozenge, \bullet$ (A), $\blacktriangle$: state points considered in Fig. 2.




FIG. 2: (a) ^3He concentration profile $X(l) = 1 - Q_l$ and (b) SOP m_l profile for a VBEG film of width $L = 60$ for $K = 0.5J$, $\Delta_1/J = -3$, and $\Delta_2/J = \Delta_t/J \simeq 0.61$ corresponding to the state points $\blacklozenge$, $\bullet$, and $\blacktriangle$ indicated in Fig. 1.




FIG. 3: Color online. Dimensionless scaling function $\vartheta(y = tL) = f_C L^3$ for the VBEG model calculated within MF theory along the paths of fixed concentration of ^3He shown in Fig. 1. Dots indicate the corresponding onset temperature $T_s(y)$ of superfluidity at the λ -line.




FIG. 4: The adjusted scaling function $\bar{\vartheta}(y)$ (see main text) for the VBEG model within MFT compared with the corresponding experimental curve [4] obtained along the path of fixed tricritical concentration $X_t \approx 0.672$ of ^3He .